



Cambridge IGCSE™

CAMBRIDGE INTERNATIONAL MATHEMATICS

0607/05

Paper 5 Investigation (Core)

For examination from 2025

MARK SCHEME B

Maximum Mark: 40

Specimen

This document has **8** pages. Any blank pages are indicated.

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptions for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptions for the question
- the specific skills defined in the mark scheme or in the generic level descriptions for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptions.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptions in mind.

Mathematics-Specific Marking Principles

- 1 Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
- 2 Unless specified in the question, non-integer answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
- 3 Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
- 4 Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
- 5 Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 A or B mark for the misread.
- 6 Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

MARK SCHEME NOTES

The following notes are intended to help with understanding of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Anything in the mark scheme which is in square brackets [...] is not required for the mark to be earned, but if present it must be correct.

When a part of a question has two or more ‘method’ steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. The notation ‘dep’ is used to indicate that a particular M or B mark is dependent on an earlier mark in the scheme.

Types of mark

- M** Method mark, awarded for a valid method applied to the problem.
- A** Accuracy mark, given for a correct answer or intermediate step correctly obtained. For accuracy marks to be given, the associated Method mark must be earned or implied.
- B** Mark for a correct result or statement independent of Method marks.
- C** Communication mark, for clear mathematical communication and reasoning.

Abbreviations

awrt	answers which round to
cao	correct answer only
dep	dependent on the previous mark(s)
FT	follow through after error
isw	ignore subsequent working (after correct answer obtained)
nfw	not from wrong working
oe	or equivalent
SC	special case
soi	seen or implied

Question	Answer	Marks	Partial Marks																																							
1	4 16 25	1																																								
2(a)(i)	81	1																																								
2(a)(ii)	1600	1																																								
2(b)	$81 + 1600 = 1681 = 41^2$	2	B1 for 1681 seen																																							
3	At least three differences in the 2nd or the 3rd column or $c = \sqrt{a^2 + b^2}$ or $a^2 + b^2 = c^2$ used	C1																																								
	<table border="1"> <thead> <tr> <th><i>a</i></th> <th><i>b</i></th> <th><i>c</i></th> </tr> </thead> <tbody> <tr><td>3</td><td>4</td><td>5</td></tr> <tr><td>5</td><td>12</td><td>13</td></tr> <tr><td>7</td><td>24</td><td>25</td></tr> <tr><td>9</td><td>40</td><td>41</td></tr> <tr><td>11</td><td>60</td><td>61</td></tr> <tr><td>13</td><td>84</td><td>85</td></tr> <tr><td>15</td><td>112</td><td>113</td></tr> <tr><td>17</td><td>144</td><td>145</td></tr> <tr><td>19</td><td>180</td><td>181</td></tr> <tr><td>21</td><td>220</td><td>221</td></tr> <tr><td>23</td><td>264</td><td>265</td></tr> <tr><td>25</td><td>312</td><td>313</td></tr> </tbody> </table>	<i>a</i>	<i>b</i>	<i>c</i>	3	4	5	5	12	13	7	24	25	9	40	41	11	60	61	13	84	85	15	112	113	17	144	145	19	180	181	21	220	221	23	264	265	25	312	313	5	B1 for first column AND In middle and last columns B4 for all 6 cells correct, or B3 for 4 cells correct, or B2 for 2 cells correct, or B1 for 1 cell correct.
<i>a</i>	<i>b</i>	<i>c</i>																																								
3	4	5																																								
5	12	13																																								
7	24	25																																								
9	40	41																																								
11	60	61																																								
13	84	85																																								
15	112	113																																								
17	144	145																																								
19	180	181																																								
21	220	221																																								
23	264	265																																								
25	312	313																																								

Question	Answer	Marks	Partial Marks
4	Link: (3, 4, 5) and (5, 12, 13) or (7, 24, 25) and (25, 312, 313)	C1	
	(3, 4, 12, 13) (7, 24, 312, 313)	2	B1 for each
5(a)	$7^2 + 14^2 + 14^2 = 441$ $21^2 = 441$	2	B1 for each statement
5(b)	$k^2a^2 + k^2b^2 + k^2c^2 = k^2d^2$	M1	
	Evidence of dividing by k^2 or $k^2(a^2 + b^2 + c^2)$ seen leading to $a^2 + b^2 + c^2 = d^2$	A1	
5(c)(i)	7	1	
5(c)(ii)	Division by 7 seen	C1	
	(1, 2, 2, 3)	1	
5(d)(i)	6	1	
5(d)(ii)	(2, 6, 9, 11)	1	
5(d)(iii)	$2^2 + 6^2 + 9^2 = 121$ $11^2 = 121$	2	FT (4, 12, 18, 22) or (6, 18, 27, 33) B1 for each statement or B1FT <i>their part (ii)</i> for one correct statement
6(a)(i)	$\frac{2^2 + 3^2 - 1}{2}$ or $\frac{2^2 + 3^2 + 1}{2}$	C1	
	(2, 3, 6, 7)	2	B1 for 6 or 7 correctly placed
6(a)(ii)	$\frac{2^2 + b^2 + 1}{2} = 43$	C1	
	(2, 9, 42, 43)	2	B1 for 9 or 42 correctly placed

Question	Answer	Marks	Partial Marks																							
6(b)(i)	At least 3 differences in one of the columns b , c or d	C1																								
	<table border="1"> <thead> <tr> <th>a</th> <th>b</th> <th>c</th> <th>d</th> </tr> </thead> <tbody> <tr> <td>2</td> <td>3</td> <td>6</td> <td>7</td> </tr> <tr> <td>2</td> <td>5</td> <td>14</td> <td>15</td> </tr> <tr> <td>2</td> <td>7</td> <td>26</td> <td>27</td> </tr> <tr> <td>2</td> <td>9</td> <td>42</td> <td>43</td> </tr> <tr> <td>2</td> <td>11</td> <td>62</td> <td>63</td> </tr> </tbody> </table>	a	b	c	d	2	3	6	7	2	5	14	15	2	7	26	27	2	9	42	43	2	11	62	63	2
a	b	c	d																							
2	3	6	7																							
2	5	14	15																							
2	7	26	27																							
2	9	42	43																							
2	11	62	63																							
6(b)(ii)	$d = c + 1$ oe	1																								
6(c)	Substituting a and b both even in $\frac{a^2 + b^2 - 1}{2}$ or $\frac{a^2 + b^2 + 1}{2}$	M1																								
	Correct answer or 'not an integer' oe	A1																								
6(d)(i)	$\frac{a^2 + b^2 + 1}{2} = 23$ leading to $a^2 + b^2 + 1 = 46$ or $\frac{45 + 1}{2} = 23$ or $c = 22$ and $\frac{a^2 + b^2 - 1}{2} = 22$ leading to $a^2 + b^2 - 1 = 44$ or $c = 22$ and $\frac{45 - 1}{2} = 22$	1																								
6(d)(ii)	(3, 6, 22, 23)	B2	B1 for 3 and 6 correctly placed B1 for 22 correctly placed																							

BLANK PAGE